arXiv:gr-qc/9908024v1 7 Aug 1999

Geometric Background for Thermal Field Theories

Massimo Blasone[♭♮], Yuan-Xing Gui[♭♯] and Francis Vendrell[♭†]

[♭] *Blackett Laboratory, Imperial College, Prince Consort Road, London SW7 2BZ, United Kingdom*
[♮] *Dipartimento di Fisica, Università di Salerno and INFN, I-84100 Salerno, Italy*
[♯] *Department of Physics, Dalian University of Technology, Dalian 116024, China*
[†] *Département de physique théorique, Ecole de physique, Université de Genève, CH-1211 Genève 4, Switzerland*

(February 7, 2008)

We study a new spacetime which is shown to be the general geometrical background for Thermal Field Theories at equilibrium. The different formalisms of Thermal Field Theory are unified in a simple way in this spacetime. The set of time-paths used in the Path Ordered Method is interpreted in geometrical terms.

I. INTRODUCTION

Thermal Field Theory (TFT) [1–3] is the generic name denoting Quantum Field Theories (QFT) at finite temperature, such as the Matsubara or Imaginary Time (IT) formalism [1–3], Thermo Field Dynamics (TFD) [1,4], and the Path Ordered Method (POM), which comprises the Closed Time Path (CTP) formalism [1–3] as a particular case. These distinct approaches result from different efforts to introduce a temperature within the framework of quantum field theory.

The IT formalism exploits the analogy of the temperature with imaginary time in calculating the partition function, and within this formalism the two-point Green's function is given by the Matsubara propagator. In contrast, the POM formalism and TFD both deal with real time. These two last approaches lead to the same matrix structure for the propagator in equilibrium, although their founding ideas are different. In the POM formalism, the temperature is introduced by adding a pure imaginary number to the real time and by choosing a special path in the complex-time plane. The propagators are calculated by taking functional derivatives of a path integral over fields, the so-called finite-temperature generating functional. On the other hand, in TFD, the field algebra is doubled and the temperature is contained explicitly in the resulting "vacuum" state, which is then referred to as the "thermal vacuum". The propagators are expressed in this last formalism as expectation values of time-ordered products of quantum fields with respect to the thermal vacuum.

An important and interesting question is whether such different formalisms have some roots in common and if their features can be understood in a deeper way so that they appear unified. A clue to an answer may be found in the well-known discovery of Hawking [5] that temperature arises in a quantum theory as a result of a non-trivial background endowed with event-horizon(s), in this case a black-hole spacetime. Rindler spacetime, the spacetime of an accelerated observer, was also shown later to exhibit thermal features [6,7]. There is also a flat background with a non-trivial structure which exhibits thermal features, the so-called η-ξ spacetime [8–10], which has besides revealed itself useful for discussing properties of black holes [11]. We believe that η-ξ spacetime is the generic example of a spacetime exhibiting thermal properties in the sense that fields and states in this spacetime look everywhere as if they were immersed in a thermal bath contained in a Minkowski background [8,9].

η-ξ spacetime is a flat complex manifold with topology $S^1 \times R^3$. Its Lorentzian section is made up of four copies of Minkowski spacetime glued together along some of their past or future null hyperplanes infinities. In Kruskal-like coordinates, which cover the entire η-ξ spacetime, the metric is singular on these hyperplanes. For this reason they are *formally* called "event-horizons". Their existence leads to the doubling of the degrees of freedom of the fields. The vacuum propagator on the Lorentzian section corresponds to the real-time thermal matrix propagator. In the Euclidean section of Kruskal coordinates, the time coordinate is periodic, so that the propagator is equal to the Matsubara propagator there.

In real-time TFTs, there is a freedom in the parameterization of the thermal matrix propagator. In the POM formalism, this parameterization is related to the choice of the path in the complex-time plane going from $t = 0$ to $t = -i\beta$ which is not unique [12] (see Fig. 1). In TFD, different parameterizations of the Bogoliubov thermal matrix are possible. It is important to stress that although the choice of the parameterization is irrelevant in the thermal equilibrium case since it does not affect physical quantities in real-time TFTs, it nevertheless plays a role in the non-equilibrium case, where the choice of a closed time path is the only useful one [14–16].

In this paper we show the relevance of other complex sections of η-ξ spacetime than the Lorentzian and the Euclidean ones. All possible time paths used in the POM formalism, including the CTP formalism, and the different parameterizations of the Bogoliubov matrices of TFD are interpreted at a purely geometric level in these complex sections. The geometric picture for TFTs is consequently enlarged. We also show how the different formalisms of TFT unify themselves in a very natural way in this framework. The generalization introduced here could be useful in order to extend the geometrical picture of η-ξ spacetime to systems out of thermal equilibrium. We consider in detail the case of a scalar field, the generalization to fermions being straightforward as shown in Ref. [10].

The paper is organized as follows: In Section II, we define the η-ξ spacetime in general and then we study its Euclidean and Lorentzian sections, and their complex extensions. In Section III, we consider explicitly a quantum boson field in η-ξ spacetime. Section IV is devoted to the study of the relationships between η-ξ spacetime and TFTs, and to the unification of TFTs in the framework of η-ξ spacetime. Section V contains a discussion of some other features of the extended Lorentzian section. Finally, Section VI is devoted to conclusions.

II. η-ξ SPACETIME

η-ξ spacetime [8,9] is a four-dimensional complex manifold defined by the line element

$$ds^2 = \frac{-d\eta^2 + d\xi^2}{\alpha^2\,(\xi^2 - \eta^2)} + dy^2 + dz^2, \tag{1}$$

where $\alpha \equiv 2\pi/\beta$ is a real constant and $(\eta, \xi, y, z) \in \mathbb{C}$. We shall use the symbol $\xi^\mu \equiv (\eta, \xi, y, z)$ to denote as a whole the set of η-ξ coordinates, but for simplicity we shall actually drop the index μ when no confusion with the space-like coordinate is possible.

A. Euclidean section

The Euclidean section of η-ξ spacetime is obtained by assuming that $(\sigma, \xi, y, z) \in \mathbb{R}$ where $\eta \equiv i\sigma$. The metric in this section is given from Eq. (1) by

$$ds^2 = \frac{d\sigma^2 + d\xi^2}{\alpha^2\,(\sigma^2 + \xi^2)} + dy^2 + dz^2. \tag{2}$$

By use of the transformation

$$\begin{cases} \sigma &= (1/\alpha)\, \exp(\alpha x) \sin(\alpha\tau)\,, \\ \xi &= (1/\alpha)\, \exp(\alpha x) \cos(\alpha\tau)\,, \end{cases} \tag{3}$$

the metric becomes that of the cylindrical Euclidean flat spacetime,

$$ds^2 = d\tau^2 + dx^2 + dy^2 + dz^2, \tag{4}$$

where the time τ has a periodic structure, i.e. $\tau \equiv \tau + \beta$.

B. Lorentzian section

In the Lorentzian section the metric is given by Eq. (1) where $(\eta, \xi, y, z) \in \mathbb{R}$. This metric is singular on the two hyperplanes $\eta = \pm\xi$ which shall be called the "event-horizons". They divide η-ξ spacetime into four regions denoted by R_{I}, R_{II}, R_{III} and R_{IV} (see Fig. 2).

In the first two regions, one defines two sets of tortoise-like coordinates $x_{I,II} \in \mathbb{R}^4$ by $x_{I,II} = (t_{I,II}, x_{I,II}, y, z)$ where

$$\text{in } R_I\text{:} \quad \begin{cases} \eta &= +(1/\alpha)\, \exp(\alpha x_I) \sinh(\alpha t_I)\,, \\ \xi &= +(1/\alpha)\, \exp(\alpha x_I) \cosh(\alpha t_I)\,, \end{cases} \tag{5}$$

$$\text{in } R_{II}\text{:} \quad \begin{cases} \eta &= -(1/\alpha)\, \exp(\alpha x_{II}) \sinh(\alpha t_{II})\,, \\ \xi &= -(1/\alpha)\, \exp(\alpha x_{II}) \cosh(\alpha t_{II})\,. \end{cases} \tag{6}$$

Similar transformations can be defined to cover R_{III} and R_{IV} (see Refs. [8,11]) but it shall be sufficient for our purposes to consider only the first two regions. The metric given in Eq. (1) becomes the Minkowski metric in these regions and in the new coordinates,

$$ds^2 = -dt^2_{I,II} + dx^2_{I,II} + dy^2 + dz^2. \tag{7}$$

Regions I to IV are copies of the Minkowski spacetime glued together along the "event-horizons" making up the Lorentzian section of η-ξ spacetime.

Although Eqs. (5) and (6) are formally Rindler transformations [6], the $(t_{I,II}, x_{I,II}, y, z)$ coordinates should not be confused with Rindler coordinates since in those coordinates the metric is the Minkowski one. An observer whose motion is described for example by $(x_I(t_I), y_I(t_I), z(t_I)) = (x_0, y_0, z_0)$ is subjected to an inertial motion, and thus cannot be identified with an accelerated observer as in the Rindler case. In Eqs. (5) and (6), the role of the inertial and non-inertial coordinates are actually reversed with respect to the Rindler case [6,8].

We now study the analytical properties of the transformations given in Eqs. (5) and (6). If $\xi^\pm = \eta \pm \xi$, we rewrite them in the form

$$\begin{cases} \xi^+_>(t_I, x_I) &= +(1/\alpha) \exp\left[+\alpha\,(t_I + x_I)\right], \\ \xi^-_<(t_I, x_I) &= -(1/\alpha) \exp\left[-\alpha\,(t_I - x_I)\right], \end{cases} \tag{8}$$

$$\begin{cases} \xi^+_<(t_{II}, x_{II}) &= -(1/\alpha) \exp\left[+\alpha\,(t_{II} + x_{II})\right], \\ \xi^-_>(t_{II}, x_{II}) &= +(1/\alpha) \exp\left[-\alpha\,(t_{II} - x_{II})\right], \end{cases} \tag{9}$$

where we have added the subscripts $<$ and $>$ to the variables $\xi^\pm$ to indicates their ranges, i.e. one has $\xi^\pm_> > 0$ and $\xi^\pm_< < 0$. The reciprocals of these transformations are

$$\begin{cases} t_{I}(\xi^+_>,\xi^-_<) &= \dfrac{1}{2\alpha}\ln\left(-\dfrac{\xi^+_>}{\xi^-_<}\right), \\ x_{I}(\xi^+_>,\xi^-_<) &= \dfrac{1}{2\alpha}\ln\left(-\alpha^2\xi^+_>\xi^-_<\right), \end{cases} \tag{10}$$

$$\begin{cases} t_{II}(\xi^+_<,\xi^-_>) &= \dfrac{1}{2\alpha}\ln\left(-\dfrac{\xi^+_<}{\xi^-_>}\right), \\ x_{II}(\xi^+_<,\xi^-_>) &= \dfrac{1}{2\alpha}\ln\left(-\alpha^2\xi^+_<\xi^-_>\right). \end{cases} \tag{11}$$

Equations (10) and (11) are defined in regions R_{I} and R_{II} respectively. We now would like to extend analytically these expressions to obtain the functions $t_{I}(\xi)$, $t_{II}(\xi)$ and $x_{I}(\xi)$, $x_{II}(\xi)$ defined in $R_{I}\cup R_{II}$. This amounts to extend these expressions from positive or negatives values of $\xi^\pm$ to their negative or positive values respectively. In order to do this, we choose to analytically extend the expressions above in the *lower* half-planes of both the ξ^+ and ξ^- complex planes for reasons which will become clear below. In other words we assume that $-\pi \leq \arg\xi^\pm < \pi$, or equivalently that the cuts in the $\xi^\pm$ complex planes are given by $\mathbb{R}_- + i0^+$.

It is not possible to perform the analytic extensions with respect to the two variables $\xi^\pm$ *at once*, otherwise an erroneous result would be obtained. To fix the ideas, we choose to perform the extension first in the ξ^+ variable and then in ξ^- (the choice of the opposite order gives the same result for our particular purposes). If $\xi^\pm_<$ is the analytic continuation of $\xi^\pm_>$ from negative to positive values, one has

$$\ln\left(-\xi^\pm_<\right) = \ln\left(+\xi^\pm_>\right) + i\pi, \tag{12}$$

$$\ln\left(+\xi^\pm_>\right) = \ln\left(-\xi^\pm_>\right) - i\pi. \tag{13}$$

This implies

$$\ln\left(-\frac{\xi^+_<}{\xi^-_>}\right) = \ln\left(-\frac{\xi^+_>}{\xi^-_<}\right) + i2\pi, \tag{14}$$

$$\ln\left(-\alpha^2\xi^+_<\xi^-_>\right) = \ln\left(-\alpha^2\xi^+_>\xi^-_<\right), \tag{15}$$

which means that these expressions are the analytic continuations of each others. In consequence, by using the variable ξ one obtains in $R_{I}\cup R_{II}$,

$$\begin{cases} t_{II}(\xi) &= t_{I}(\xi) + i\,\beta/2, \\ x_{II}(\xi) &= x_{I}(\xi). \end{cases} \tag{16}$$

C. Extended Lorentzian section

We now consider a class of complex sections of η-ξ spacetime obtained from the Lorentzian section by shifting the Minkowski time coordinates in the imaginary direction *only* in region R_{II},

$$\begin{array}{ll} \text{in } R_{I}\cup R_{III}\cup R_{IV}: & t_q \to t_{q_\delta} = t_q, \\ \text{in } R_{II}: & t_{II} \to t_{II_\delta} = t_{II} + i\beta\delta, \end{array} \tag{17}$$

where $\delta \in [-1/2, 1/2]$. The set of these sections shall be called "extended Lorentzian section." We denote by R_{II_δ} the image of the region R_{II} through this shift. In R_{II_δ} the η-ξ coordinates become complex variables and are transformed according to $(\eta,\xi) \to (\eta_\delta, \xi_\delta)$ where, from Eq. (17),

$$\begin{cases} \eta_\delta &= -(1/\alpha)\,\exp\left(\alpha x_{II}\right)\sinh\left[\alpha\left(t_{II} + i\beta\delta\right)\right], \\ \xi_\delta &= -(1/\alpha)\,\exp\left(\alpha x_{II}\right)\cosh\left[\alpha\left(t_{II} + i\beta\delta\right)\right]. \end{cases} \tag{18}$$

In terms of the real η-ξ variables, we have

$$\begin{cases} \eta_\delta &= +\eta\,\cos\left(2\pi\delta\right) + i\xi\,\sin\left(2\pi\delta\right), \\ i\xi_\delta &= -\eta\,\sin\left(2\pi\delta\right) + i\xi\,\cos\left(2\pi\delta\right), \end{cases} \tag{19}$$

or equivalently

$$\begin{cases} \xi^+_\delta &= \exp\left(+i2\pi\delta\right)\,\xi^+, \\ \xi^-_\delta &= \exp\left(-i2\pi\delta\right)\,\xi^-. \end{cases} \tag{20}$$

The time shift induces thus a rotation in the $(\eta, i\xi)$ plane of R_{II}. The metric becomes in terms of the rotated coordinates

$$ds^2 = \frac{-d\eta_\delta^2 + d\xi_\delta^2}{\alpha^2\left(\xi_\delta^2 - \eta_\delta^2\right)} + dy^2 + dz^2, \tag{21}$$

and is thus unchanged by the time shift which is therefore an isometry of the four-dimensional complex η-ξ spacetime. The equations (16) become after the time shift

$$\begin{cases} t_{II_\delta}(\xi_\delta) &= t_{I}(\xi_\delta) + i\,\beta\left(1/2 + \delta\right), \\ x_{II_\delta}(\xi_\delta) &= x_{I}(\xi_\delta). \end{cases} \tag{22}$$

III. FIELDS IN η-ξ SPACETIME

We now consider a free scalar field in the η-ξ spacetime (for the case of a fermion field see Ref. [10]). The "global" scalar field in η-ξ coordinates shall be denoted by $\Phi(\xi)$. It satisfies to the Klein-Gordon equation

$$(\Box + m^2)\Phi(\xi) = 0, \tag{23}$$

where $\Box = \nabla_\mu\nabla^\mu$ if ∇_μ is the covariant derivative. The scalar product of two fields is given by

$$(\Phi_1, \Phi_2) = -i\int_\Sigma d\Sigma\,\left\|g(\xi)\right\|^{1/2}\,\Phi_1(\xi)n^\nu\,\overleftrightarrow{\partial}_\nu\,\Phi_2^*(\xi), \tag{24}$$

where g is the determinant of the metric $g^{\mu\nu}$, Σ is any space-like surface and n^ν an orthonormal vector to this surface.

A. Euclidean section

In the Euclidean section we have $\Phi = \Phi(\sigma, \xi, y, z)$, and the field in the t-x coordinates defined in Eq. (3) shall be denoted by $\phi = \phi(\tau, x, y, z)$. These two fields are related by

$$\phi(\tau, x, y, z) = \Phi(\sigma(\tau, x), \xi(\tau, x), y, z). \tag{25}$$

Because of the periodic nature of the time τ and by imposing single valuedness we have necessarily

$$\phi(\tau, x, y, z) = \phi(\tau + \beta, x, y, z). \tag{26}$$

B. Lorentzian section

In the Lorentzian section, as we have seen, we have four different regions, each of them being a complete Minkowski spacetime. Since we are interested only in regions R_I and R_{II}, we shall consider the quantum field over these two regions only. Our aim is to find an expansion for the global field Φ in the joining $R_I \cup R_{II}$.

We start by defining the "local" fields $\phi^I(x_I)$ and $\phi^{II}(x_{II})$ by

$$\Phi(\xi) = \begin{cases} \phi^I(x_I(\xi)), & \text{when } \xi \in R_I, \\ \phi^{II}(x_{II}(\xi)), & \text{when } \xi \in R_{II}. \end{cases} \tag{27}$$

They have support in R_I and R_{II} respectively. By choosing the particular surface $\eta = a\xi$ where a is a constant satisfying $-1 < a < 1$, one shows from Eq. (24) that the global scalar product is given by

$$(\Phi_1, \Phi_2) = < \phi_1^I, \phi_2^I > + < \phi_1^{II}, \phi_2^{II} >, \tag{28}$$

where $<,>$ is the local scalar product in Minkowski spacetime

$$< \phi_1, \phi_2 > = -i \int_{\mathbb{R}^3} d^3x\, \phi_1(x) \overleftrightarrow{\partial}_t \phi_2^*(x). \tag{29}$$

In η-ξ spacetime covered by t-x coordinates given in Eqs. (5) and (6), the mode solutions of the Klein-Gordon equation are just local plane waves restricted to a given region. They are given by

$$u_{\mathbf{k}}(x_I) = (4\pi\omega_{\mathbf{k}})^{-\frac{1}{2}}\ e^{i\left(-\omega_{\mathbf{k}} t_I + \mathbf{k}\cdot\mathbf{x}_I\right)}, \tag{30}$$

$$v_{\mathbf{k}}(x_{II}) = (4\pi\omega_{\mathbf{k}})^{-\frac{1}{2}}\ e^{i\left(+\omega_{\mathbf{k}} t_{II} + \mathbf{k}\cdot\mathbf{x}_{II}\right)}, \tag{31}$$

where $\omega_{\mathbf{k}} = \sqrt{\mathbf{k}^2 + m^2}$. From these Minkowski modes, one defines the two wave functions $U_{\mathbf{k}}(\xi)$ and $V_{\mathbf{k}}(\xi)$ with support in R_I and R_{II} respectively by

$$U_{\mathbf{k}}(\xi) = \begin{cases} u_{\mathbf{k}}(x_I(\xi)), & \text{when } \xi \in R_I, \\ 0, & \text{when } \xi \in R_{II}, \end{cases} \tag{32}$$

$$V_{\mathbf{k}}(\xi) = \begin{cases} 0, & \text{when } \xi \in R_I, \\ v_{\mathbf{k}}(x_{II}(\xi)), & \text{when } \xi \in R_{II}. \end{cases} \tag{33}$$

Their power spectrum with respect to the momenta conjugated to ξ^+ and ξ^- contains negative contributions, which are furthermore not bounded by below. Consequently, the sets of functions $\left\{U_{\mathbf{k}}(\xi), U^*_{-\mathbf{k}}(\xi)\right\}_{\mathbf{k}\in\mathbb{R}^3}$ and $\left\{V_{\mathbf{k}}(\xi), V^*_{-\mathbf{k}}(\xi)\right\}_{\mathbf{k}\in\mathbb{R}^3}$ defined on R_I and R_{II} respectively are both over-complete since the same energy contribution (i.e. momentum contribution conjugate to η) can appear twice in these sets. In other words, the energy spectrum of $U_{\mathbf{k}}$ and $U^*_{-\mathbf{k}}$ overlap, and so do the ones of $V_{\mathbf{k}}$ and $V^*_{-\mathbf{k}}$. These sets can thus not be used as a basis in their respective regions, and the joining of these sets is clearly not a basis in $R_I \cup R_{II}$.

To construct a basis in $R_I \cup R_{II}$, we could solve the Klein-Gordon equations in η-ξ coordinates to obtain the field modes in these coordinates. However, the Bogoliubov transformations resulting from this basis choice are rather complicated. So instead of doing this, we shall construct from the wave functions $u_{\mathbf{k}}(x_I(\xi))$ and $v^*_{-\mathbf{k}}(x_{II}(\xi))$ basis elements having positive energy spectrum.

We shall demand these basis elements to be analytical functions in the lower complex planes of ξ^+ and ξ^-, so that their spectrum contain only positive contributions of the momenta conjugate with respect to ξ^+ and ξ^-. In consequence, they shall have positive energy spectra.

We thus extend analytically the two wave functions $u_{\mathbf{k}}(x_I(\xi))$ and $v^*_{-\mathbf{k}}(x_{II}(\xi))$ in the lower complex planes of ξ^+ and ξ^- (as in Section II B, the cut in the complex planes is given by $\mathbb{R}_- + i0^+$). By applying formula (16) we get directly

$$u_{\mathbf{k}}(x_I(\xi)) = e^{-\frac{\beta}{2}\omega_{\mathbf{k}}}\, v^*_{-\mathbf{k}}(x_{II}(\xi)), \tag{34}$$

$$v_{\mathbf{k}}(x_{II}(\xi)) = e^{-\frac{\beta}{2}\omega_{\mathbf{k}}}\, u^*_{-\mathbf{k}}(x_I(\xi)). \tag{35}$$

The expressions on the right and left hand sides of these last equations are analytic continuations of each others. In this way we are led to introduce the two normalized linear combinations

$$\begin{cases} F_{\mathbf{k}}(\xi) \ = \ (1 - f_{\mathbf{k}})^{-\frac{1}{2}} \left[U_{\mathbf{k}}(\xi) + f_{\mathbf{k}}^{\frac{1}{2}}\, V^*_{-\mathbf{k}}(\xi)\right], \\ \widetilde{F}_{\mathbf{k}}(\xi) \ = \ (1 - f_{\mathbf{k}})^{-\frac{1}{2}} \left[V_{\mathbf{k}}(\xi) + f_{\mathbf{k}}^{\frac{1}{2}}\, U^*_{-\mathbf{k}}(\xi)\right], \end{cases} \tag{36}$$

where $f_{\mathbf{k}} = e^{-\beta\omega_{\mathbf{k}}}$, and where $U_{\mathbf{k}}(\xi)$ and $V_{\mathbf{k}}(\xi)$ are defined in Eqs. (32) and (33). These wave functions are still solutions of the Klein-Gordon equation. They are analytical in $R_I \cup R_{II}$ and in particular at the origin $\xi^+ = \xi^- = 0$. Since they are analytical complex functions in the lower complex planes of ξ^+ and ξ^-, their spectrum has only positive energy contributions. The set $\{F_{\mathbf{k}}, F^*_{-\mathbf{k}}, \widetilde{F}_{\mathbf{k}}, \widetilde{F}^*_{-\mathbf{k}}\}_{\mathbf{k}\in\mathbb{R}^3}$ is thus complete but not overcomplete over the joining $R_I \cup R_{II}$. Furthermore it is an orthogonal set since

$$\begin{array}{rcccl} (F_{\mathbf{k}}, F_{\mathbf{p}}) & = & (\widetilde{F}^*_{\mathbf{k}}, \widetilde{F}^*_{\mathbf{p}}) & = & +\delta^3(\mathbf{k}-\mathbf{p}), \\ (F^*_{\mathbf{k}}, F^*_{\mathbf{p}}) & = & (\widetilde{F}_{\mathbf{k}}, \widetilde{F}_{\mathbf{p}}) & = & -\delta^3(\mathbf{k}-\mathbf{p}), \end{array} \tag{37}$$

with all the other scalar products vanishing.

On one hand, the local scalar fields can be expanded in the Minkowski modes given in Eqs. (30) and (31),

$$\phi^I(x_I) = \int d^3\mathbf{k} \left[a^I_{\mathbf{k}}\, u_{\mathbf{k}}(x_I) + a^{I\,\dagger}_{\mathbf{k}}\, u^*_{\mathbf{k}}(x_I) \right], \tag{38}$$

$$\phi^{II}(x_{II}) = \int d^3\mathbf{k} \left[a^{II}_{\mathbf{k}}\, v_{\mathbf{k}}(x_{II}) + a^{II\,\dagger}_{\mathbf{k}}\, v^*_{\mathbf{k}}(x_{II}) \right]. \tag{39}$$

And on the other hand, the global scalar field can be expanded in terms of the "global" modes given in Eqs. (36) as

$$\Phi(\xi) = \int d^3\mathbf{k} \Big[b_{\mathbf{k}}\, F_{\mathbf{k}}(\xi) + b^\dagger_{\mathbf{k}}\, F^*_{-\mathbf{k}}(\xi) + \tilde{b}_{\mathbf{k}}\, \widetilde{F}_{\mathbf{k}}(\xi) + \tilde{b}^\dagger_{\mathbf{k}}\, \widetilde{F}^*_{-\mathbf{k}}(\xi) \Big]. \tag{40}$$

These three expansions define the local and global creation and annihilation operators, which are related by Bogoliubov transformations. To obtain these, we recall the definition (27) relating the local and global fields and we use the field expansions (38), (39) and (40). We obtain

$$\left\{ \begin{array}{rcl} b_{\mathbf{k}} & = & a^I_{\mathbf{k}} \cosh\theta_{\mathbf{k}} - a^{II\,\dagger}_{\mathbf{k}} \sinh\theta_{\mathbf{k}}, \\ \tilde{b}_{\mathbf{k}} & = & a^{II}_{\mathbf{k}} \cosh\theta_{\mathbf{k}} - a^{I\,\dagger}_{\mathbf{k}} \sinh\theta_{\mathbf{k}}, \end{array} \right. \tag{41}$$

where $\sinh^2\theta_{\mathbf{k}} = n(\omega_{\mathbf{k}}) = (e^{\beta\omega_{\mathbf{k}}} - 1)^{-1}$.

C. Extended Lorentzian section

By following the above procedure, we shall now construct a set of positive energy modes defined in the extended Lorentzian section introduced in Section II C. We start by considering region R_{II}, where the set $\{v_{\mathbf{k}}(x_{II}), v^*_{-\mathbf{k}}(x_{II})\}_{\mathbf{k}\in\mathbb{R}^3}$ is a plane wave basis. One has

$$v_{\mathbf{k}}(x_{II}) = (4\pi\omega_{\mathbf{k}})^{-\frac{1}{2}}\, e^{i\left(+\omega_{\mathbf{k}} t_{II} + \mathbf{k}\cdot\mathbf{x}_{II}\right)}, \tag{42}$$

$$v^*_{-\mathbf{k}}(x_{II}) = (4\pi\omega_{\mathbf{k}})^{-\frac{1}{2}}\, e^{i\left(-\omega_{\mathbf{k}} t_{II} + \mathbf{k}\cdot\mathbf{x}_{II}\right)}. \tag{43}$$

Under the time shift, Eq. (17), this set is transformed into $\{v_{\mathbf{k}}(x_{II_\delta}), v^\sharp_{-\mathbf{k}}(x_{II_\delta})\}_{\mathbf{k}\in\mathbb{R}^3}$, where we have replaced the symbol $*$ by $\sharp$ because $v^\sharp_{-\mathbf{k}}(x_{II_\delta})$ is no longer the complex conjugate of $v_{\mathbf{k}}(x_{II_\delta})$. Indeed, one has

$$v_{\mathbf{k}}(x_{II_\delta}(x_{II})) = e^{-\beta\omega_{\mathbf{k}}\delta}\, v_{\mathbf{k}}(x_{II}), \tag{44}$$

$$v^\sharp_{-\mathbf{k}}(x_{II_\delta}(x_{II})) = e^{+\beta\omega_{\mathbf{k}}\delta}\, v^*_{-\mathbf{k}}(x_{II}). \tag{45}$$

The complex conjugation and the time shift are not commutative. We notice that $v^\sharp_{-\mathbf{k}}(x_{II_\delta}(x_{II}))$ can be obtained from $v_{\mathbf{k}}(x_{II_\delta}(x_{II}))$ by complex conjugation *and* by the replacement $\delta \to -\delta$. This rule actually defines the $\sharp$-conjugation.

In the same way as in Eq. (27) one defines

$$\Phi(\xi) = \left\{ \begin{array}{ll} \phi^I(x_I(\xi)), & \text{when } \xi \in R_I, \\ \phi^{II_\delta}(x_{II_\delta}(\xi)), & \text{when } \xi \in R_{II_\delta}. \end{array} \right. \tag{46}$$

The global scalar product, Eq. (28), is modified and becomes in $R_I \cup R_{II_\delta}$,

$$(\Phi_1, \Phi_2)_\delta = <\phi^I_1, \phi^I_2> + <\phi^{II_\delta}_1, \phi^{II_\delta}_2>_\delta, \tag{47}$$

where the local Minkowski scalar product $< \,,\, >_\delta$ in region R_{II_δ} is given by

$$<\phi_1, \phi_2>_\delta = -i \int_{\mathbb{R}^3} d^3x_{II_\delta}\, \phi_1(x_{II_\delta}) \overleftrightarrow{\partial}_t\, \phi^\sharp_2(x_{II_\delta}). \tag{48}$$

Equations (34), (35), (44) and (45) imply

$$u_{\mathbf{k}}(x_I(\xi_\delta)) = e^{-\beta\omega_{\mathbf{k}}(1/2+\delta)}\, v^\sharp_{-\mathbf{k}}(x_{II_\delta}(\xi_\delta)), \tag{49}$$

$$v_{\mathbf{k}}(x_{II_\delta}(\xi_\delta)) = e^{-\beta\omega_{\mathbf{k}}(1/2+\delta)}\, u^\sharp_{-\mathbf{k}}(x_I(\xi_\delta)). \tag{50}$$

Consequently one also has

$$u^\sharp_{\mathbf{k}}(x_I(\xi_\delta)) = e^{-\beta\omega_{\mathbf{k}}(1/2-\delta)}\, v_{-\mathbf{k}}(x_{II_\delta}(\xi_\delta)), \tag{51}$$

$$v^\sharp_{\mathbf{k}}(x_{II_\delta}(\xi_\delta)) = e^{-\beta\omega_{\mathbf{k}}(1/2-\delta)}\, u_{-\mathbf{k}}(x_I(\xi_\delta)). \tag{52}$$

The expressions on the left and right hand sides of these equations are thus analytic continuations of each others. If we now define

$$U_{\mathbf{k}}(\xi_\delta) = \left\{ \begin{array}{ll} u_{\mathbf{k}}(x_I(\xi_\delta)), & \text{when } \xi_\delta \in R_I, \\ 0, & \text{when } \xi_\delta \in R_{II_\delta}, \end{array} \right. \tag{53}$$

$$V_{\mathbf{k}}(\xi_\delta) = \left\{ \begin{array}{ll} 0, & \text{when } \xi_\delta \in R_I, \\ v_{\mathbf{k}}(x_{II_\delta}(\xi_\delta)), & \text{when } \xi_\delta \in R_{II_\delta}, \end{array} \right. \tag{54}$$

the global modes in the extended Lorentzian section $R_I \cup R_{II_\delta}$ are given by

$$\left\{ \begin{array}{rcl} G_{\mathbf{k}}(\xi_\delta) & = & (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left[U_{\mathbf{k}}(\xi_\delta) + f_{\mathbf{k}}^{\frac{1}{2}+\delta}\, V^\sharp_{-\mathbf{k}}(\xi_\delta) \right], \\ \widetilde{G}_{\mathbf{k}}(\xi_\delta) & = & (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left[V_{\mathbf{k}}(\xi_\delta) + f_{\mathbf{k}}^{\frac{1}{2}+\delta}\, U^\sharp_{-\mathbf{k}}(\xi_\delta) \right], \end{array} \right. \tag{55}$$

and

$$\left\{ \begin{array}{lcl} G^{\sharp}_{\mathbf{k}}(\xi_\delta) &=& (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left[U^{\sharp}_{\mathbf{k}}(\xi_\delta) + f_{\mathbf{k}}^{\frac{1}{2}-\delta}\, V_{-\mathbf{k}}(\xi_\delta)\right], \\ \widetilde{G}^{\sharp}_{\mathbf{k}}(\xi_\delta) &=& (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left[V^{\sharp}_{\mathbf{k}}(\xi_\delta) + f_{\mathbf{k}}^{\frac{1}{2}-\delta}\, U_{-\mathbf{k}}(\xi_\delta)\right], \end{array} \right. \tag{56}$$

where $f_{\mathbf{k}} = e^{-\beta\omega_{\mathbf{k}}}$. From Eqs. (49) to (52), we see that the global modes are analytic in $R_{_I} \cup R_{_{II_\delta}}$, in particular at the origin $\xi^+_\delta = \xi^-_\delta = 0$. Since they are analytical complex functions on the lower complex planes of ξ^+_δ and ξ^-_δ, their spectrum has only positive energy contributions. When $\delta = 0$, the above combinations consistently reduce to the expressions of Eq. (36).

We emphasize that the global modes $G^*_{\mathbf{k}}$ and $\widetilde{G}^*_{\mathbf{k}}$ are *not* analytic in the extended Lorentzian section, contrary to the the non-hermitian combinations $G^{\sharp}_{\mathbf{k}}$ and $\widetilde{G}^{\sharp}_{\mathbf{k}}$. Non-hermitian conjugation operations such as our sharp conjugation $\sharp$ are actually common in TFT, see for example Ref. [18] for a formally similar situation. In Ref. [19] it is shown the necessity of the so-called Osterwalder-Schrader reflection as opposed to the hermiticity property in Euclidean field theories even when the temperature vanishes.

The set $\{G_{\mathbf{k}}, \widetilde{G}_{\mathbf{k}}, G^{\sharp}_{\mathbf{k}}, \widetilde{G}^{\sharp}_{\mathbf{k}}\}_{\mathbf{k}\in\mathbb{R}^3}$ is thus complete over $R_{_I} \cup R_{_{II_\delta}}$. It is furthermore an orthogonal set since

$$\begin{array}{lclcl} (G_{\mathbf{k}}, G_{\mathbf{p}})_\delta &=& (\widetilde{G}^{\sharp}_{\mathbf{k}}, \widetilde{G}^{\sharp}_{\mathbf{p}})_\delta &=& +\delta^3(\mathbf{k}-\mathbf{p}), \\ (G^{\sharp}_{\mathbf{k}}, G^{\sharp}_{\mathbf{p}})_\delta &=& (\widetilde{G}_{\mathbf{k}}, \widetilde{G}_{\mathbf{p}})_\delta &=& -\delta^3(\mathbf{k}-\mathbf{p}), \end{array} \tag{57}$$

where all the other scalar products vanish.

On one hand, the expansions of the local fields in the Minkowski modes over regions $R_{_I}$ and $R_{_{II_\delta}}$ are given respectively by

$$\phi^{_I}(x_{_I}) = \int d^3\mathbf{k} \left[a^{_I}_{\mathbf{k}}\, u_{\mathbf{k}}(x_{_I}) + a^{_I\,\dagger}_{\mathbf{k}}\, u^*_{\mathbf{k}}(x_{_I})\right], \tag{58}$$

$$\phi^{_{II_\delta}}(x_{_{II_\delta}}) = \int d^3\mathbf{k} \left[a^{_{II_\delta}}_{\mathbf{k}}\, v_{\mathbf{k}}(x_{_{II_\delta}}) + a^{_{II_\delta}\,\dagger}_{\mathbf{k}}\, v^*_{\mathbf{k}}(x_{_{II_\delta}})\right]. \tag{59}$$

On the other hand, the expansion of the global field in the G modes over the joining $R_{_I} \cup R_{_{II_\delta}}$ is

$$\begin{aligned} \Phi(\xi_\delta) = \int d^3\mathbf{k} \Big[& c_{\mathbf{k}}\, G_{\mathbf{k}}(\xi_\delta) + c^{\sharp}_{\mathbf{k}}\, G^{\sharp}_{\mathbf{k}}(\xi_\delta) \\ & + \tilde{c}_{\mathbf{k}}\, \tilde{G}_{-\mathbf{k}}(\xi_\delta) + \tilde{c}^{\sharp}_{\mathbf{k}}\, \tilde{G}^{\sharp}_{-\mathbf{k}}(\xi_\delta)\Big]. \end{aligned} \tag{60}$$

From these last expansions and by using Eqs. (55) and (56), one finds the Bogoliubov transformations,

$$\left\{ \begin{array}{lcl} c_{\mathbf{k}} &=& (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left(a^{_I}_{\mathbf{k}} - f_{\mathbf{k}}^{\frac{1}{2}-\delta}\, a^{_{II_\delta}\,\dagger}_{\mathbf{k}}\right), \\ \tilde{c}_{\mathbf{k}} &=& (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left(a^{_{II_\delta}}_{\mathbf{k}} - f_{\mathbf{k}}^{\frac{1}{2}-\delta}\, a^{_I\,\dagger}_{\mathbf{k}}\right), \end{array} \right. \tag{61}$$

and their $\sharp$-conjugate equivalent,

$$\left\{ \begin{array}{lcl} c^{\sharp}_{\mathbf{k}} &=& (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left(a^{_I\,\dagger}_{\mathbf{k}} - f_{\mathbf{k}}^{\frac{1}{2}+\delta}\, a^{_{II_\delta}}_{\mathbf{k}}\right), \\ \tilde{c}^{\sharp}_{\mathbf{k}} &=& (1-f_{\mathbf{k}})^{-\frac{1}{2}} \left(a^{_{II_\delta}\,\dagger}_{\mathbf{k}} - f_{\mathbf{k}}^{\frac{1}{2}+\delta}\, a^{_I}_{\mathbf{k}}\right). \end{array} \right. \tag{62}$$

Again the transformations given in Eq. (41) are recovered when $\delta = 0$.

IV. THERMAL FIELD THEORIES IN η-ξ SPACETIME

We now show the equivalence of Quantum Field Theory in η-ξ spacetime with Thermal Field Theories by considering the scalar field. We shall see that in several sections of η-ξ spacetime QFT naturally reproduces the known formalisms of TFTs insofar as the correct thermal Green functions are recovered there.

On one hand, we shall show that in the Euclidean section of η-ξ spacetime QFT corresponds to the imaginary time formalism, and that the Green functions are the Matsubara Green functions in this section. On the other hand, we shall show that in the extended Lorentzian section QFT reproduces the two known formalisms of TFTs with real time, namely the POM formalism and TFD. Furthermore we shall fix the parameter δ of the extended Lorentzian section with respect to the parameter σ of the POM formalism and of TFD. The most general thermal matrix propagator shall be obtained in the framework of η-ξ spacetime. We shall need below the covariant Lagrangian for the scalar field with a source term J. It is given by

$$\mathcal{L}[\Phi, \mathrm{J}] = \sqrt{-g}\left(\frac{1}{2}\,\partial_\mu\Phi\partial_\nu\Phi + \frac{m^2}{2}\,\Phi^2 - V(\Phi) - \mathrm{J}\Phi\right). \tag{63}$$

A. Imaginary time formalism

The generating functional for the Green functions in the Euclidean section of η-ξ spacetime is [8]

$$Z_E[J] = N \int [d\Phi] \exp\left\{-\int d\sigma d\xi dy dz\ \mathcal{L}_{\sigma,\xi}[\Phi, \mathrm{J}]\right\}, \tag{64}$$

where

$$\begin{aligned} \mathcal{L}_{\sigma,\xi}[\Phi, \mathrm{J}] = & \frac{1}{2}\,\left[(\partial_\sigma\Phi)^2 + (\partial_\xi\Phi)^2\right] \\ & + \frac{1}{\alpha^2\,(\sigma^2+\xi^2)}\left\{\frac{1}{2}\,(\nabla_\perp\Phi)^2 + \frac{m^2}{2}\,\Phi^2 + V(\Phi) - \mathrm{J}\Phi\right\}. \end{aligned} \tag{65}$$

We now perform the change of coordinates given in Eq. (3) in the generating functional. In the τ-x coordinates, the sum over fields is taken over all periodic fields satisfying to the periodic constraint in Eq. (26),

$$Z_E[J] = N \int\limits_{\beta-periodic} [d\phi] \exp\left\{ -\int_0^\beta d\tau \int_{\mathbb{R}^3} dxdydz\, \mathcal{L}_{\tau,x}[\phi, J] \right\} \tag{66}$$

where $J(\tau, x, y, z) = \mathrm{J}(\sigma, \xi, y, z)$ and

$$\mathcal{L}_{\tau,x}[\phi, J] = \frac{1}{2}\left[(\partial_\tau \phi)^2 + (\nabla\phi)^2 + m^2\,\phi^2 \right] + V(\phi) - J\phi. \tag{67}$$

By differentiation this last equation with respect to the source J we obtain the Matsubara propagator whose Fourier transforms is

$$G_\beta(\mathbf{k}, \omega_n) = \frac{1}{\omega_n^2 + \mathbf{k}^2 + m^2}, \tag{68}$$

where the Matsubara frequencies ω_n are given for bosons by $\omega_n = 2\pi n/\beta$ ($n \in \mathbb{N}$).

B. Real time formalism: Path Ordered Method

Let us now consider the extended Lorentzian section of η-ξ spacetime. The generating functional is given by [8]

$$Z[J] = \mathcal{N}\!\int [d\Phi] \exp\left\{ i \int d\eta_\delta d\xi_\delta dydz\, \mathcal{L}_{\eta_\delta,\xi_\delta}[\Phi, \mathrm{J}] \right\}, \tag{69}$$

where

$$\mathcal{L}_{\eta_\delta,\xi_\delta}[\Phi, \mathrm{J}] = \frac{1}{2}\left[\left(\partial_{\eta_\delta}\Phi\right)^2 - \left(\partial_{\xi_\delta}\Phi\right)^2 \right] \tag{70}$$

$$+ \frac{1}{\alpha^2\left(\eta_\delta^2 - \xi_\delta^2\right)} \left\{ -\frac{1}{2}\left(\nabla_\perp \Phi\right)^2 - \frac{m^2}{2}\Phi^2 - V(\Phi) + \mathrm{J}\Phi \right\}.$$

Since we shall be interested only in the propagators whose spacetime arguments belong only to the joining $R_{I} \cup R_{II_\delta}$, we can set the source to zero in regions R_{III} and R_{IV},

$$\mathrm{J}(x) = 0, \qquad\qquad \text{when } x \in R_{III} \cup R_{IV}. \tag{71}$$

Then the expression in Eq. (69) reduces to

$$Z[J] = \mathcal{N} \int [d\Phi] \exp\left\{ i \int_{R_I \cup R_{II_\delta}} d\eta_\delta d\xi_\delta dydz\ \mathcal{L}_{\eta_\delta,\xi_\delta}[\Phi, \mathrm{J}] \right\}. \tag{72}$$

We now express the fields in regions R_I and R_{II_δ} in terms of the local Minkowskian coordinates by using the transformations given in Eq. (18). We obtain

$$Z[J] = \mathcal{N} \int [d\phi] \exp\left\{ i \int dt_I dx_I dydz\ \mathcal{L}_{t,x}[\phi, J] \right. \tag{73}$$

$$\left. + i \int dt_{II_\delta} dx_{II_\delta} dydz\ \mathcal{L}_{t,x}[\phi, J] \right\},$$

where the integration is taken over the Minkowski spacetime, ϕ is the local field and where

$$\mathcal{L}_{t,x}[\phi, J] = \frac{1}{2}\left[(\partial_t \phi)^2 - (\nabla\phi)^2 - m^2\,\phi^2 \right] - V(\phi) + J\phi. \tag{74}$$

We now use the relations (22) and obtain

$$Z[J] = \mathcal{N} \int [d\phi] \exp\left\{ i \int dtdxdydz \right. \tag{75}$$

$$\left. \times \left[\mathcal{L}_{t,x}[\phi, J]\,(t, x) - \mathcal{L}_{t,x}[\phi, J]\,(t + i\beta\delta, x) \right] \right\},$$

where in the last step we have dropped the subscript I and where we have taken into account the fact that the direction of time in R_{II_δ} is opposite to the one in R_I, resulting in a minus sign in the second integration.

We now consider the expression for the generating functional as given in the POM formalism [2]. It is given by

$$Z_{\text{POM}}[J] = \mathcal{N}' \int [d\phi] \exp\left\{ i \int_C dtdxdydz\ \mathcal{L}_{t,x}[\phi, J] \right\}, \tag{76}$$

where the time path C is shown in Fig. 1. The contribution from the vertical parts of the contour may be included in the normalization factor $\mathcal{N}'$ and neglected when calculating the real-time Green functions. The generating functionals in Eq. (75) and (76) can then be identified provided that

$$\delta = \sigma - 1/2. \tag{77}$$

We thus see that the type of time path in the POM formalism is related directly to the "rotation angle" between the two regions R_I and R_{II_δ} of η-ξ spacetime. In the free field case, from the above generating functionals we obtain the well-known thermal matrix propagator [1–3],

$$\begin{aligned}
D_{11}(k) &= \frac{i}{k^2 - m^2 + i0^+} + 2\pi\, n(k_0)\, \delta(k^2 - m^2), \\
D_{22}(k) &= D_{11}^*(k), \\
D_{12}(k) &= e^{\sigma\beta k_0}\left[n(k_0) + \theta(-k_0) \right] 2\pi\, \delta(k^2 - m^2), \\
D_{21}(k) &= e^{-\sigma\beta k_0}\left[n(k_0) + \theta(k_0) \right] 2\pi\, \delta(k^2 - m^2),
\end{aligned} \tag{78}$$

where $n(k_0) = (e^{\beta|k_0|} - 1)^{-1}$. The parameter σ appears explicitly only in the off-diagonal components of the matrix propagator.

C. Real time formalism: Thermo Field Dynamics

Another formalism for real-time TFT is Thermo Field Dynamics [4,13]. In this approach a central role is played by the Bogoliubov transformation, relating the zero-temperature annihilation and creation operators with the thermal ones. In TFD the field algebra is doubled and one then considers two commuting field operators ϕ and $\tilde{\phi}$ given by

$$\phi(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}(2\omega_k)^{\frac{1}{2}}} \left[a_{\mathbf{k}} e^{i(-\omega_k t + \mathbf{kx})} + a^\dagger_{\mathbf{k}} e^{i(\omega_k t - \mathbf{kx})}\right], \tag{79}$$

$$\tilde{\phi}(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}(2\omega_k)^{\frac{1}{2}}} \left[\tilde{a}_{\mathbf{k}} e^{i(\omega_k t - \mathbf{kx})} + \tilde{a}^\dagger_{\mathbf{k}} e^{i(-\omega_k t + \mathbf{kx})}\right]. \tag{80}$$

In the original formulation of Takahashi and Umezawa [4], the thermal Bogoliubov transformation is given by

$$\left\{ \begin{array}{rcl} a_{\mathbf{k}}(\theta) &=& a_{\mathbf{k}} \cosh\theta_{\mathbf{k}} - \tilde{a}^\dagger_{\mathbf{k}} \sinh\theta_{\mathbf{k}}, \\ \tilde{a}_{\mathbf{k}}(\theta) &=& \tilde{a}_{\mathbf{k}} \cosh\theta_{\mathbf{k}} - a^\dagger_{\mathbf{k}} \sinh\theta_{\mathbf{k}}, \end{array} \right. \tag{81}$$

where $\sinh^2\theta_{\mathbf{k}} = n(\omega_k)$. These operators annihilate the "thermal vacuum" $|0(\beta)\rangle$. Thermal averages are calculated as expectation values with respect to $|0(\beta)\rangle$.

The form of the thermal Bogoliubov matrix is however not unique. It is possible to generalize the above transformation to a non-hermitian superposition of the form [13,14]

$$\left\{ \begin{array}{rcl} \gamma_{\mathbf{k}} &=& (1 - f_{\mathbf{k}})^{-\frac{1}{2}} \left(a_{\mathbf{k}} - f^{1-\sigma}_{\mathbf{k}}\, \tilde{a}^\dagger_{\mathbf{k}}\right), \\ \tilde{\gamma}_{\mathbf{k}} &=& (1 - f_{\mathbf{k}})^{-\frac{1}{2}} \left(\tilde{a}_{\mathbf{k}} - f^{1-\sigma}_{\mathbf{k}}\, a^\dagger_{\mathbf{k}}\right), \end{array} \right. \tag{82}$$

where $f_{\mathbf{k}} = e^{-\beta\omega_{\mathbf{k}}}$. The non-hermiticity property of this last transformation implies that the canonical conjugates of γ and $\tilde{\gamma}$ are not $\gamma^\dagger$ and $\tilde{\gamma}^\dagger$, but are rather the combinations

$$\left\{ \begin{array}{rcl} \gamma^\sharp_{\mathbf{k}} &=& (1 - f_{\mathbf{k}})^{-\frac{1}{2}} \left(a^\dagger_{\mathbf{k}} - f^{\sigma}_{\mathbf{k}}\, \tilde{a}_{\mathbf{k}}\right), \\ \tilde{\gamma}^\sharp_{\mathbf{k}} &=& (1 - f_{\mathbf{k}})^{-\frac{1}{2}} \left(\tilde{a}^\dagger_{\mathbf{k}} - f^{\sigma}_{\mathbf{k}}\, a_{\mathbf{k}}\right), \end{array} \right. \tag{83}$$

which give the correct canonical commutators, $[\gamma_{\mathbf{k}}, \gamma^\sharp_{\mathbf{p}}] = \delta^3(\mathbf{k} - \mathbf{p})$, etc. Here the $\sharp$ conjugation corresponds to the usual hermitian conjugation $\dagger$ *together* with the replacement $\sigma \to 1 - \sigma$. The hermitian representation in Eq. (81) is recovered when $\sigma = 1/2$.

Thermal averages are now expressed as [13,14]

$$\langle A \rangle = \frac{{}_L\langle 0(\beta)|A|0(\beta)\rangle_R}{{}_L\langle 0(\beta)|0(\beta)\rangle_R}, \tag{84}$$

where A is an observable, and where $|0(\beta)\rangle_R$ and ${}_L\langle 0(\beta)|$ are the left and right vacuum states defined by

$$\left. \begin{array}{c} \gamma \\ \tilde{\gamma} \end{array} \right\} |0(\beta)\rangle_R \,=\, 0 \,=\, {}_L\langle 0(\beta)| \left\{ \begin{array}{c} \gamma^\sharp \\ \tilde{\gamma}^\sharp \end{array} \right. \tag{85}$$

The thermal propagator for a scalar field is calculated in TFD as follows:

$$D^{(ab)}(x, y) = \langle T\left[\phi^a(x)\phi^b(y)^\dagger\right]\rangle, \tag{86}$$

where

$$\phi^a \equiv \begin{pmatrix} \phi \\ \tilde{\phi} \end{pmatrix}. \tag{87}$$

This propagator coincides with the one of Eqs. (78), as can be easily checked by using the definitions given above. The connection of TFD with the geometrical picture of η-ξ spacetime is immediate if we make the identification

$$\begin{pmatrix} \phi \\ \tilde{\phi} \end{pmatrix} \equiv \begin{pmatrix} \phi^{I} \\ \phi^{II_\delta} \end{pmatrix} \tag{88}$$

under the constraint of Eq. (77) (see also Eq. (46)). Then the Bogoliubov transformations in Eqs. (61), (62) and those in Eqs. (82), (83) are identical.

V. OTHER FEATURES OF THE EXTENDED LORENTZIAN SECTION

In this Section we discuss other features of the η-ξ spacetime, which are influenced by the rotation of Eq. (19). Along the lines of Ref. [20], we consider the analytic continuation of the imaginary time thermal propagator to real times in the context of η-ξ spacetime. In Ref. [20] it was shown that the geometric structure of η-ξ spacetime plays a crucial role in obtaining the matrix real-time propagator from the Matsubara one.

To see how this works, it is sufficient to consider the simple case of a massless free scalar field in two-dimensions. In the Euclidean section of η-ξ spacetime, the equation for the propagator is

$$\left(\frac{\partial^2}{\partial\sigma^2} + \frac{\partial^2}{\partial\xi^2}\right) D_I(A - A') = -(-g_E)^{-\frac{1}{2}} \delta^2(A - A'), \tag{89}$$

where (A, A') denotes a couple of points, g_E is the determinant of the metric in the Euclidean section, and where D_I is the imaginary time thermal propagator. Now we continue Eq. (89) to the extended Lorentzian section. This is achieved by first replacing σ by $i\eta$ and then performing the rotation of Eq. (19). If D is the real time propagator, we have

$$\left(-\frac{\partial^2}{\partial\eta_\delta^2} + \frac{\partial^2}{\partial\xi_\delta^2}\right) D(A - A') = -(-g_L)^{-\frac{1}{2}} \delta^2(A - A'), \tag{90}$$

where g_L stands for the determinant of the metric in the Lorentzian section. Because of the presence of different disconnected regions in the Lorentzian section, the propagator has a matrix structure, since now the points A and

A' can belong either to region R_I or R_{II_δ} (R_{III} and R_{IV} are excluded since they are space-like with respect to R_I and R_{II_δ}). The delta function with complex arguments A and A' is defined to be zero when the points belong to different regions.

In Minkowski coordinates, Eq. (90) reads

$$\left(-\frac{\partial^2}{\partial t^2}+\frac{\partial^2}{\partial x^2}\right)D(A-A')=-\delta^2_C(A-A'). \qquad (91)$$

The delta function containing the time variable in the last equation needs to be defined on an appropriate path since the time variable is complex. This path coincides with the POM time path of Fig. 1 when the identification of Eq. (77) is made. By use of Eq. (22) and by following the procedure of Ref. [20] we obtain, for example for the component D_{12}, the equation

$$\left(-\frac{\partial^2}{\partial t^2}+\frac{\partial^2}{\partial x^2}\right)D(t-t'+i\sigma\beta,x-x') \\ =-\delta_C(t-t'+i\sigma\beta)\,\delta(x-x'), \qquad (92)$$

which has the D_{12} propagator given in Eq. (78) as solution.

Let us finally consider the *tilde conjugation* in the context of η-ξ spacetime. The tilde conjugation rules are postulated in TFD in order to connect the physical and the tilde operators. Due to the geometrical structure of η-ξ spacetime, these rules are there seen as coordinate transformations. This was first discussed in Ref. [21]. We extend here the result to the extended Lorentzian section of η-ξ spacetime.

Let us recall the tilde rules as defined in TFD [4,13] (we restrict for simplicity to bosonic operators):

$$\begin{array}{lll} (AB)\tilde{} = \tilde{A}\tilde{B}, & (C_1A+C_2B)\tilde{} = C_1^*\tilde{A}+C_2^*\tilde{B}, \\ \left(\tilde{A}\right)\tilde{} = A, & \left(\tilde{A}\right)^\dagger = \left(A^\dagger\right)\tilde{}, \end{array} \qquad (93)$$

where A,B are operators and C_1,C_2 are c-numbers. In order to reproduce this operation in the extended Lorentzian section of η-ξ spacetime, let us first introduce the following M operation as defined in Ref. [21].

$$M\,\Phi(\eta,\xi)\,M^{-1}\equiv\Phi(-\eta,-\xi). \qquad (94)$$

When the field is expressed in Minkowskian coordinates the M operation reads as

$$M\,\phi(t,x)\,M^{-1}=\phi(t-i\beta/2,x). \qquad (95)$$

The M operation is anti-linear, since it induces a time inversion together with the shift $t\to t-i\beta/2$, see Ref. [21]. This is clear when we consider its action on the conjugate momentum $\pi(t,x)\equiv\partial_t\phi^\dagger(t,x)$,

$$M\,\pi(t,x)\,M^{-1}=-\pi(t-i\beta/2,x). \qquad (96)$$

Next we perform a rotation by an angle δ transforming the η,ξ coordinates according to Eqs. (19) and the field becomes then

$$R_\delta\,\Phi(\eta,\xi)\,R_\delta^{-1}\equiv\Phi(\eta_\delta,\xi_\delta). \qquad (97)$$

Finally we introduce a δ conjugation operation, which is similar to a charge conjugation, by

$$C_\delta\,\phi(t,x)\,C_\delta^{-1}\equiv\phi^\sharp(t,x), \qquad (98)$$

where the change $\delta\to-\delta$ (or equivalently $\sigma\to1-\sigma$) has to be performed together with usual charge conjugation.

The combined action of the three operations results in the tilde conjugation. By defining for simplicity the notation $G_\delta\equiv C_\delta\,R_\delta\,M$, we get

$$\begin{aligned} G_\delta\,\phi(t,x)\,G_\delta^{-1} &= \phi^\sharp(t-i\sigma\beta,x) \\ &= \phi^\dagger(t-i(1-\sigma)\beta,x). \end{aligned}$$

We then get by omitting the space dependence for simplicity

$$\begin{aligned} G_\delta\,\phi_1(t)\,\phi_2(t')\,G_\delta^{-1} &= \phi_1^\sharp(t-i\sigma\beta)\,\phi_2^\sharp(t'-i\sigma\beta), \\ G_\delta\left[G_\delta\,\phi(t)\,G_\delta^{-1}\right]G_\delta^{-1} &= \phi(t), \\ G_\delta\left[B_1\,\phi_1(t)+B_2\,\phi_2(t')\right]G_\delta^{-1} & \\ &= B_1^*\,\phi_1^\sharp(t-i\sigma\beta)+B_2^*\,\phi_2^\sharp(t'-i\sigma\beta), \\ G_\delta\left[\phi^\dagger(t)\right]G_\delta^{-1} &= \left[G_\delta\,\phi(t)\,G_\delta^{-1}\right]^\dagger. \end{aligned} \qquad (99)$$

The c-numbers are conjugated since the M operation is anti-linear. The second of the above relations is derived as follows

$$\begin{aligned} & G_\delta\left[G_\delta\,\phi(t)\,G_\delta^{-1}\right]G_\delta^{-1} \\ &= C_\delta\,R_\delta\,M\,\phi^\dagger(t-i(1-\sigma)\beta)\,M^{-1}\,R_\delta^{-1}\,C_\delta^{-1} \\ &= C_\delta\,\phi^\dagger(t-i(1-\sigma)\beta-i\sigma\beta)\,C_\delta^{-1} \\ &= \phi(t-i\beta)=\phi(t). \end{aligned} \qquad (100)$$

The tilde rules of Eq. (93) are thus reproduced.

VI. DISCUSSION AND CONCLUSIONS

We have discussed a section of η-ξ spacetime which represent the general geometric background for real-time thermal field theories at equilibrium. This section is complex and can be regarded as an extension of the usual Lorentzian section of η-ξ spacetime by means of a rotation of region R_{II} with respect to R_I in the complex η-ξ spacetime. In terms of Minkowski coordinates the rotation is equivalent to a constant time shift, leaving the metric invariant.

The angle between the two regions turns out to be related to the σ parameter of the time path as used in the POM formalism. It also reproduces the parameter present in the Bogoliubov thermal matrix of TFD, when the relation between modes belonging to different regions is considered. The general form of the thermal matrix propagator containing the parameter σ has been obtained by use of this particular geometrical background. Finally, we have discussed the analytic continuation of imaginary time propagator to real time matrix propagator and the tilde rule in the context of η-ξ spacetime.

In the geometrical background of η-ξ spacetime, it is possible to understand the differences between the various formalisms of TFT in a simple way. In particular, with regards to the real-time methods, i.e. the POM formalism and TFD, the geometric picture is the following. In the Lorentzian section of η-ξ spacetime we have two different regions $R_{_I}$ and $R_{_{II_\delta}}$ over which the field is defined. For a global observer this field is a free field. However when one does restrict to one of the two regions (say $R_{_I}$), temperature arise as a consequence of the loss of information (increase in entropy) about the other region. In order to calculate the propagator, one needs then to compare the fields defined in different regions. This can be done essentially in two ways:

1) By analytically continuing the field $\phi^{_{II_\delta}}(x_{_{II_\delta}})$ defined in region $R_{_{II_\delta}}$ to region $R_{_I}$. Then the time argument gets shifted by $i\beta(1/2+\delta)$, as described in Section III. One thus ends up with *one field* and *two possible time arguments*, which can be either t or $t - i\beta(1/2+\delta)$. The generating functional defined in η-ξ spacetime by following this procedure turns out to be the same of the one defined in the POM formalism.

2) One can attach the information about the region to the field operator rather than putting it in the time argument. Thus the identification $\phi^{_I}(x) \equiv \phi(x)$ and $\phi^{_{II_\delta}}(x) \equiv \tilde{\phi}(x)$ can be made and one obtains the formalism of TFD, which consists of *two commuting field operators* and *a single time argument.*

Of course, the two pictures give the same physics, i.e. the same propagator, since they are only different "viewpoints" of local observers in the context of η-ξ spacetime. It is an interesting question to ask if such an equivalence can be extended for situation which are out of thermal equilibrium. Work is in progress along this direction.

ACKNOWLEDGEMENTS

We thank Dr. T. S. Evans for many useful discussions. M. B. thanks MURST, INFN and ESF, Y. X. G. the Natural National Science Foundation of China and the Royal Society of UK, and F. V. the Swiss National Science Foundation.

[1] A. Das, *Finite Temperature Field Theory* (World Scientific, 1997).

[2] M. Le Bellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, 1996).

[3] N. P. Landsman and Ch. G. van Weert, *Phys. Rep.* **145** (1987) 141.

[4] Y. Takahashi and H. Umezawa, *Collective Phenomena* **2** (1975) 55; reprinted in *Int. J. Mod. Phys.* **B10** (1996) 1755; H. Umezawa, H. Matsumoto and M. Tachiki, *Thermo Field Dynamics and Condensed States*, (North-Holland Publ. Co., Amsterdam, 1982).

[5] S. W. Hawking, *Commun. Math. Phys.* **43** (1975) 199.

[6] W. Rindler, *Am. J. Phys.* **34** (1966) 1174.

[7] N. D. Birrell and P. C. W. Davies, *Quantum Field in Curved Space*, (Cambridge University Press), 1982.

[8] Y. X. Gui, *Phys. Rev.* **D 42** (1990) 1988.

[9] Y. X. Gui, *Phys. Rev.* **D46** (1992) 1869.

[10] Y. X. Gui, *Phys. Rev.* **D45** (1992) 697.

[11] F. Vendrell, gr-qc/9906008; M. E. Ortiz and F. Vendrell, *Phys. Rev.* **D59** (1999) 084005.

[12] A. J. Niemi and G. Semenoff, *Ann. Phys.* **152** (1984) 105.

[13] H. Umezawa, *Advanced Field Theory: Micro, Macro and Thermal Physics* (American Institute of Physics, New York, 1993).

[14] P. A. Henning, *Phys. Rep.* **253** (1995) 235.

[15] P. A. Henning, M. Graf and F. Matthäus, *Physica* **A 182** (1992) 489.

[16] T. S. Evans, I. Hardman, H. Umezawa and Y. Yamanaka, *J. Math. Phys.* **33** (1992) 370.

[17] Y. X. Gui, *Sci. Sin.* **A 36** (1993) 561.

[18] M. Blasone, T. S. Evans, D. Steer and G. Vitiello, *J. Phys.* **A 32** (1999) 1185.

[19] K. Osterwalder and R. Schrader, *Comm. Math. Phys.* **31** (1973) 83; **42** (1975) 281.

[20] J. Zuo and Y. X. Gui, hep-ph/9803235.

[21] J. Zuo and Y. X. Gui, *J. Phys.* **A 28** (1995) 4907.



FIG. 1. The time path used in POM. The parameter σ ranges from the value $\sigma = 0$ (Closed Time Path) to $\sigma = 1$.



FIG. 2. Lorentzian section of η-ξ spacetime: the solid lines represent the singularities at $\xi^2 - \eta^2 = 0$. On the straight lines time is constant, while on the hyperbolas the Minkowski coordinate x is constant. Note that time flows in opposite directions in regions I and II.